FACTS ABOUT ALKANES

Physical State Carbon Count : State 1-4 Gases 5 - 15 Liquid 15 - more Solid

Preparation of Alkanes

1. From unsaturated Hydrocarbons in the presence of catalyst 'Ni' or 'Pt'

$$CH_2=CH_2+H_2\xrightarrow{Ni}CH_3-CH_3$$

2. By Wurtz Reaction:

$$CH_3 - Br + Na + Br - CH_3 \xrightarrow{Dry} CH_3 - CH_3 + 2NaBr$$

3. By the Reduction of Aldehydes and Ketones

$$\begin{array}{c} 0 \\ R-\ddot{\mathbb{C}}-H+4[H] \xrightarrow{Zn.Hg/conc.HCL} R-CH_3 + H_2O \end{array}$$

Grignard's Reagent

$$R-Mg-X + H_2O \longrightarrow R-H + HO-Mg-X$$



Density of alkanes is less than water therefore they float over it.

Solubility

Alkanes do not dissolve in water. They form a layer on top of water. However alkanes dissolve in non-polar organic solvents like Toulene, Benzene



Example of Alkanes

Methane




Ethane



Propane



C - in sp3 Hybridisation



General Formula :- C.H2n+2

Uses

Lighter alkanes are used in natural gas.

Propane and Butane are used in LPG cylinders



Boiling

Boiling point depends on Vander waal forces.



More Carbon atoms



High Vander waal forces



High boiling point



Melting

Melting point depends on packing of compound.



More branching



Close packing



High melting point



ALKENES


Physical State

	Carbon Count	State
	1-3	Gases
ı	4-20	Liquid
ı	> 20	Solid

Acts as a Nucleophile

In organic chemistry, π -bond is considered as a nucleophile. Therefore alkenes participate in addition reactions

SP² hybridisation



Polymerisation

Ethene undergoes polymerisation and forms products like polyethene



Preparation

1. Dehydration of alcohols

A molecule of water is eliminated from an alcohol molecule by heating the alcohol in the presence of a strong mineral acid.

2. Dehydrohalogenation of alkyl halides

The dehydrohalogenation of alkyl halides, another β elimination reaction, involves the loss of a hydrogen and a halide from an alkyl halide (RX).

ethyl chloride

Dipole Moments


cis > 0. trans = 0

Melting Point

trans > cis

Boiling Point

cis > trans



If you add bromine water to alkene, it decolourises the liquid because bromine reacts with alkene, where as with alkanes it cannot react.

Isomerism




sodium chloride



H C H

ALKYNES

SP hybridisation



Physical Properties



Alkynes are gases at room temperature.

0350

192 kcal/mol energy is required to break the triple bond.



Shortest bond length is 120 Å.

Test for Alkynes

Reaction with Bromine

The alkynes react slowly with bromine water to decolourise it, and this reaction can be used to distinguish between alkenes and alkynes.

Alkenes decolourise bromine water very rapidly, but alkynes take several minues.

$$R-C \equiv C-H \xrightarrow{Br_3} C = C \xrightarrow{H} Br \xrightarrow{Br} Br \xrightarrow{I} Br \xrightarrow{I$$

Acidic Hydrogen

Order of electronegativity is

$$sp > sp^2 > sp^3$$

Due to large electronegativity of sp carbon, terminal hydrogen becomes acidic and reacts with bases and undergoes neutralization.

For Terminal Alkynes

Terminal alkynes have acidic hydrogen, therefore by reacting with CuCl in NH₄OH, acidic hydrogen is replaced with Cu giving red colour.

$$R - C \equiv C - H + CuCl \longrightarrow R - C \equiv C - Cu$$
(Red)



Uses

Alkynes don't have any commercial use. Acetylene is used in oxy – acetylene flame.



Nucleophile: Like alkenes, they also act as nucleophiles, due to presence of 2π bonds

Preparation: Alkynes are prepared by hydrolysing carbides.

