


# Equilibrium

## Multiple Choice Questions (MCQs)

**Q. 1** We know that the relationship between  $K_c$  and  $K_p$  is

$$K_p = K_c (RT)^{\Delta n}$$

What would be the value of  $\Delta n$  for the reaction?




- (a) 1                      (b) 0.5                      (c) 1.5                      (d) 2

**Ans. (d)** The relationship between  $K_p$  and  $K_c$  is

$$K_p = K_c (RT)^{\Delta n}$$

where,  $\Delta n = (\text{number of moles of gaseous products}) - (\text{number of moles of gaseous reactants})$

For the reaction,



**Q. 2** For the reaction,  $\text{H}_2(g) + \text{I}_2(g) \rightleftharpoons 2 \text{HI}(g)$ , the standard free energy is  $\Delta G^\circ > 0$ . The equilibrium constant ( $K$ ) would be

- (a)  $K = 0$                       (b)  $K > 1$                       (c)  $K = 1$                       (d)  $K < 1$

**Ans. (d)**  $\Delta G^\circ$  and  $K$  are related as

$$\Delta G^\circ = -RT \ln K_c$$


when  $G^\circ > 0$  means  $\Delta G^\circ$  is positive. This can be so only if  $\ln K_c$  is negative i.e.,  $K_c < 1$ .

**Q. 3** Which of the following is not a general characteristic of equilibria involving physical processes?

- (a) Equilibrium is possible only in a closed system at a given temperature  
 (b) All measurable properties of the system remain constant  
 (c) All the physical processes stop at equilibrium  
 (d) The opposing processes occur at the same rate and there is dynamic but stable condition

**Ans. (c)** At the stage of equilibria involving physical processes like melting of ice and freezing of water etc., process does not stop but the opposite processes *i.e.*, forward and reverse process occur with the same rate.

**Q. 4**  $\text{PCl}_5$ ,  $\text{PCl}_3$ , and  $\text{Cl}_2$  are at equilibrium at 500 K in a closed container and their concentrations are  $0.8 \times 10^{-3} \text{ mol L}^{-1}$ ,  $1.2 \times 10^{-3} \text{ mol L}^{-1}$  and  $1.2 \times 10^{-3} \text{ mol L}^{-1}$ , respectively. The value of  $K_c$  for the reaction



- (a)  $1.8 \times 10^3 \text{ mol L}^{-1}$  (b)  $1.8 \times 10^{-3}$   
(c)  $1.8 \times 10^{-3} \text{ mol}^1 \text{ L}$  (d)  $0.55 \times 10^4$

**Ans. (b)** For the reaction,  $\text{PCl}_5 \rightleftharpoons \text{PCl}_3 + \text{Cl}_2$   
At 500 K in a closed container,  $[\text{PCl}_5] = 0.8 \times 10^{-3} \text{ mol L}^{-1}$

$$[\text{PCl}_3] = 1.2 \times 10^{-3} \text{ mol L}^{-1}$$

$$[\text{Cl}_2] = 1.2 \times 10^{-3} \text{ mol L}^{-1}$$

$$K_c = \frac{[\text{PCl}_3][\text{Cl}_2]}{[\text{PCl}_5]} = \frac{(1.2 \times 10^{-3}) \times (1.2 \times 10^{-3})}{(0.8 \times 10^{-3})} = 1.8 \times 10^{-3}$$


**Q. 5** Which of the following statements is incorrect?

- (a) In equilibrium mixture of ice and water kept in perfectly insulated flask, mass of ice and water does not change with time  
(b) The intensity of red colour increases when oxalic acid is added to a solution containing iron (III) nitrate and potassium thiocyanate  
(c) On addition of catalyst the equilibrium constant value is not affected  
(d) Equilibrium constant for a reaction with negative  $\Delta H$  value decreases as the temperature increases

**Ans. (b)** In the reaction,  $\text{Fe}^{3+} + \text{SCN}^- \rightleftharpoons \text{FeSCN}^{2+}$   
(Red)

When oxalic acid is added it combines with  $\text{Fe}^{3+}$  ions, then, equilibrium shifts towards backward direction and intensity of red colour decreases.

**Q. 6** When hydrochloric acid is added to cobalt nitrate solution at room temperature, the following reaction takes place and the reaction mixture becomes blue. On cooling the mixture it becomes pink. On the basis of this information mark the correct answer.




- (a)  $\Delta H > 0$  for the reaction  
(b)  $\Delta H < 0$  for the reaction  
(c)  $\Delta H = 0$  for the reaction  
(d) The sign of  $\Delta H$  cannot be predicted on the basis of this information

**Ans. (a)** In the reaction,  
 $\underset{\text{(Pink)}}{[\text{Co}(\text{H}_2\text{O})_6]^{3+}}(aq) + 4\text{Cl}^-(aq) \rightleftharpoons \underset{\text{(Blue)}}{[\text{CoCl}_4]^{2-}}(aq) + 6\text{H}_2\text{O}(l)$


On cooling, the equilibrium shifts backward direction or on heating, the equilibrium shifts forward direction. Hence, reaction is endothermic. *i.e.*,  $\Delta H > 0$ .



**Ans. (a)** For the reaction,



For the reaction,



When, the above two reactions are added, their equilibrium constants are multiplied, thus

$$K_{a_3} = \frac{[\text{H}^+]^2 [\text{S}^{2-}]}{[\text{H}_2\text{S}]} = K_{a_1} \times K_{a_2}$$

Hence,

$$K_{a_3} = K_{a_1} \times K_{a_2}$$

**Q. 10** Acidity of  $\text{BF}_3$  can be explained on the basis of which of the following concepts?


- (a) Arrhenius concept
- (b) Bronsted Lowry concept
- (c) Lewis concept
- (d) Bronsted Lowry as well as Lewis concept

**Ans. (c)** GN Lewis in 1923 defined an acid as a species which accepts an electron pair and base which donates an electron pair. As  $\text{BF}_3$  is an electron deficient compound, hence, it is a Lewis acid.

**Q. 11** Which of the following will produce a buffer solution when mixed in equal volumes?

- (a)  $0.1 \text{ mol dm}^{-3} \text{ NH}_4\text{OH}$  and  $0.1 \text{ mol dm}^{-3} \text{ HCl}$
- (b)  $0.05 \text{ mol dm}^{-3} \text{ NH}_4\text{OH}$  and  $0.1 \text{ mol dm}^{-3} \text{ HCl}$
- (c)  $0.1 \text{ mol dm}^{-3} \text{ NH}_4\text{OH}$  and  $0.05 \text{ mol dm}^{-3} \text{ HCl}$
- (d)  $0.1 \text{ mol dm}^{-3} \text{ CH}_3\text{COONa}$  and  $0.1 \text{ mol dm}^{-3} \text{ NaOH}$

**Ans. (c)** When the concentration of  $\text{NH}_4\text{OH}$  (weak base) is higher than the strong acid ( $\text{HCl}$ ), a mixture of weak base and its conjugate acid is obtained, which acts as basic buffer.



**Q. 12** In which of the following solvents is silver chloride most soluble?

- (a)  $0.1 \text{ mol dm}^{-3} \text{ AgNO}_3$  solution
- (b)  $0.1 \text{ mol dm}^{-3} \text{ HCl}$  solution
- (c)  $\text{H}_2\text{O}$
- (d) Aqueous ammonia

**Ans. (d)** Among the given solvent,  $\text{AgCl}$  is most soluble in aqueous ammonia solution.  $\text{AgCl}$  react with aqueous ammonia to form a complex,  $[\text{Ag}(\text{NH}_3)_2]^+ \text{Cl}^-$ .

**Q. 13** What will be the value of pH of  $0.01 \text{ mol dm}^{-3} \text{ CH}_3\text{COOH}$  ( $K_a = 1.74 \times 10^{-5}$ )?

- (a) 3.4 (b) 3.6 (c) 3.9 (d) 3.0

**Ans. (a)** Given that,  $K_a = 1.74 \times 10^{-5}$

Concentration of  $\text{CH}_3\text{COOH} = 0.01 \text{ mol dm}^{-3}$

$$\begin{aligned} [\text{H}^+] &= \sqrt{K_a \cdot C} \\ &= \sqrt{1.74 \times 10^{-5} \times 0.01} = 4.17 \times 10^{-4} \\ \text{pH} &= -\log [\text{H}^+] \\ &= -\log (4.17 \times 10^{-4}) = 3.4 \end{aligned}$$

**Q. 14**  $K_a$  for  $\text{CH}_3\text{COOH}$  is  $1.8 \times 10^{-5}$  and  $K_b$  for  $\text{NH}_4\text{OH}$  is  $1.8 \times 10^{-5}$ . The pH of ammonium acetate will be

- (a) 7.005 (b) 4.75 (c) 7.0 (d) Between 6 and 7

**Ans. (c)** Given that,  $K_a$  for  $\text{CH}_3\text{COOH} = 1.8 \times 10^{-5}$

$K_b$  for  $\text{NH}_4\text{OH} = 1.8 \times 10^{-5}$

Ammonium acetate is a salt of weak acid and weak base. For such salts

$$\begin{aligned} \text{pH} &= 7 + \frac{\text{p}K_a - \text{p}K_b}{2} \\ &= 7 + \frac{[-\log 1.8 \times 10^{-5}] - [-\log 1.8 \times 10^{-5}]}{2} \\ &= 7 + \frac{4.74 - 4.74}{2} = 7.00 \end{aligned}$$


**Q. 15** Which of the following options will be correct for the stage of half completion of the reaction  $A \rightleftharpoons B$ ?

- (a)  $\Delta G^\circ = 0$  (b)  $\Delta G^\circ > 0$  (c)  $\Delta G^\circ < 0$  (d)  $\Delta G^\circ = -RT \ln K$

**Ans. (a)** As we know that

$$\Delta G^\circ = -RT \ln K$$


At the stage of half completion of the reaction,



Therefore,  $K = 1$ .

Thus,  $\Delta G^\circ = 0$

**Q. 16** On increasing the pressure, in which direction will the gas phase reaction proceed to re-establish equilibrium, is predicted by applying the Le-Chatelier's principle. Consider the reaction,



Which of the following is correct, if the total pressure at which the equilibrium is established, is increased without changing the temperature?

- (a)  $K$  will remain same  
(b)  $K$  will decrease  
(c)  $K$  will increase  
(d)  $K$  will increase initially and decrease when pressure is very high

**Ans. (a)** In the reaction,  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$

If the total pressure at which the equilibrium is established, is increased without changing the temperature,  $K$  will remain same.  $K$  changes only with change in temperature.

**Q. 17** What will be the correct order of vapour pressure of water, acetone and ether at  $30^\circ\text{C}$ ? Given that among these compounds, water has maximum boiling point and ether has minimum boiling point?


- (a) Water < ether < acetone                      (b) Water < acetone < ether  
(c) Ether < acetone < water                      (d) Acetone < ether < water

**Ans. (b)** The given compounds are


Water, acetone, ether  
(Maximum b.p.)                      (Minimum b.p.)  
Greater the boiling point, lower is the vapour pressure of the solvent. Hence, the correct order of vapour pressure will be

Water < acetone < ether.

**Q. 18** At 500 K, equilibrium constant,  $K_c$ , for the following reaction is 5.



What would be the equilibrium constant  $K_c$  for the reaction?




- (a) 0.04                      (b) 0.4                      (c) 25                      (d) 2.5

**Ans. (a)** For the reaction,  $\frac{1}{2} \text{H}_2(\text{g}) + \frac{1}{2} \text{I}_2(\text{g}) \rightleftharpoons \text{HI}(\text{g})$

$$K_c = \frac{[\text{HI}]}{[\text{H}_2]^{1/2} [\text{I}_2]^{1/2}} = 5$$

Thus, for the reaction,



$$K_{c1} = \frac{[\text{H}_2] [\text{I}_2]}{[\text{HI}]^2} = \left( \frac{1}{K_c} \right)^2 = \left( \frac{1}{5} \right)^2 = \frac{1}{25} = 0.04$$



**Q. 19** In which of the following reactions, the equilibrium remains unaffected on addition of small amount of argon at constant volume?

- (a)  $\text{H}_2(\text{g}) + \text{I}_2(\text{g}) \rightleftharpoons 2\text{HI}(\text{g})$   
(b)  $\text{PCl}_5(\text{g}) \rightleftharpoons \text{PCl}_3(\text{g}) + \text{Cl}_2(\text{g})$   
(c)  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$   
(d) The equilibrium will remain unaffected in all the three cases

### 💡 Thinking Process

At constant volume, the equilibrium remain unaffected on addition of small amount of inert gas like argon, neon, krypton, etc.

**Ans. (d)** In these reactions, at constant volume



The equilibrium constant ( $K$ ) remains unaffected on addition of inert gas in all the three cases.

## Multiple Choice Questions (More Than One Options)

- Q. 20** For the reaction  $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ , the value of  $K$  is 50 at 400 K and 1700 at 500 K. Which of the following option(s) is/are correct?
- (a) The reaction is endothermic
  - (b) The reaction is exothermic
  - (c) If  $\text{NO}_2(\text{g})$  and  $\text{N}_2\text{O}_4(\text{g})$  are mixed at 400 K at partial pressures 20 bar and 2 bar respectively, more  $\text{N}_2\text{O}_4(\text{g})$  will be formed
  - (d) The entropy of the system increases

**Ans. (a, c, d)**

For the reaction,  $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$

At 400 K,  $K = 50$

At 500 K,  $K = 1700$

- (a) As the value of  $K$  increase with increase of temperature and  $K = \frac{K_f}{K_b}$ , this means that  $K_f$  increases, i.e., forward reaction is favoured. Hence, reaction is endothermic.
- (c) Since, number of moles of gaseous products are greater than the number of moles of gaseous reactants. Thus, higher pressure favours the backward reaction, i.e., more  $\text{N}_2\text{O}_4(\text{g})$  will be obtained, if  $P_{\text{product}} > P_{\text{reactant}}$ .
- (d) As reaction is accompanied by increase in the number of moles, entropy increases.

- Q. 21** At a particular temperature and atmospheric pressure, the solid and liquid phases of a pure substance can exist in equilibrium. Which of the following term defines this temperature?

- (a) Normal melting point
- (b) Equilibrium temperature
- (c) Boiling point
- (d) Freezing point


**Ans. (a, d)**

At a particular temperature and atmospheric pressure, the solid and liquid phases of a pure substance can exist as Solid  $\rightleftharpoons$  liquid.


They exists at normal melting point or normal freezing point.

## Short Answer Type Questions

- Q. 22** The ionisation of hydrochloric acid in water is given below



Label two conjugate acid-base pairs in this ionisation.



**Note** If Bronsted acid is a strong acid then its conjugate base is a weak base and vice-versa. Generally, the conjugate acid has one extra proton and each conjugate base has one less proton.

**Q. 23** The aqueous solution of sugar does not conduct electricity. However, when sodium chloride is added to water, it conducts electricity. How will you explain this statement on the basis of ionisation and how is it affected by concentration of sodium chloride?

**Ans.** Explanation for the given statement on the basis of ionisation and effect upon the concentration of sodium chloride is given below

- (i) Sugar being a non-electrolyte does not ionise in water whereas NaCl ionises completely in water and produces  $\text{Na}^+$  and  $\text{Cl}^-$  ion which help in the conduction of electricity.
- (ii) When concentration of NaCl is increased, more  $\text{Na}^+$  and  $\text{Cl}^-$  ions will be produced. Hence, conductance or conductivity of the solution increases.

**Q. 24**  $\text{BF}_3$  does not have proton but still acts as an acid and reacts with  $\ddot{\text{N}}\text{H}_3$ . Why is it so? What type of bond is formed between the two?

**Ans.**  $\text{BF}_3$  is an electron deficient compound and hence acts as Lewis acid.  $\ddot{\text{N}}\text{H}_3$  has one lone pair which it can donate to  $\text{BF}_3$  and form a coordinate bond. Hence,  $\text{NH}_3$  acts as a Lewis base.



**Q. 25** Ionisation constant of a weak base MOH, is given expression

$$K_b = \frac{[\text{M}^+][\text{OH}^-]}{[\text{MOH}]}$$

Values of ionisation constant of some weak bases at a particular temperature are given below

Base	Dimethylamine	Urea	Pyridine	Ammonia
$K_b$	$5.4 \times 10^{-4}$	$1.3 \times 10^{-14}$	$1.77 \times 10^{-9}$	$1.77 \times 10^{-5}$

Arrange the bases in decreasing order of the extent of their ionisation at equilibrium. Which of the above base is the strongest?

**Ans.** Given that, ionisation constant of a weak base MOH

$$K_b = \frac{[\text{M}^+][\text{OH}^-]}{[\text{MOH}]}$$

Larger the ionisation constant ( $K_b$ ) of a base, greater is its ionisation and stronger the base. Hence, dimethyl amine is the strongest base.

$$K_b \text{ Dimethyl amine} > \text{ammonia} > \text{pyridine} > \text{urea}$$

$$5.4 \times 10^{-4} \quad 1.77 \times 10^{-5} \quad 1.77 \times 10^{-9} \quad 1.3 \times 10^{-14}$$


**Q. 26** Conjugate acid of a weak base is always stronger. What will be the decreasing order of basic strength of the following conjugate bases?



**Ans.** Conjugate acid of the given bases are  $\text{H}_2\text{O}$ ,  $\text{ROH}$ ,  $\text{CH}_3\text{COOH}$  and  $\text{HCl}$ . Order of their acidic strength is



Hence, order of basic strength of their conjugate bases is



**Q. 27** Arrange the following in increasing order of pH.




**Ans.** (i)  $\text{KNO}_3$  is a salt of strong acid ( $\text{HNO}_3$ ) strong base ( $\text{KOH}$ ), hence its aqueous solution is neutral;  $\text{pH} = 7$ .

(ii)  $\text{CH}_3\text{COONa}$  is a salt of weak acid ( $\text{CH}_3\text{COOH}$ ) and strong base ( $\text{NaOH}$ ), hence, its aqueous solution is basic;  $\text{pH} > 7$ .

(iii)  $\text{NH}_4\text{Cl}$  is a salt of strong acid ( $\text{HCl}$ ) and weak base ( $\text{NH}_4\text{OH}$ ) hence, its aqueous solution is acidic;  $\text{pH} < 7$ .

(iv)  $\text{C}_6\text{H}_5\text{COONH}_4$  is a salt of weak acid,  $\text{C}_6\text{H}_5\text{COOH}$  and weak base,  $\text{NH}_4\text{OH}$ . But  $\text{NH}_4\text{OH}$  is slightly stronger than  $\text{C}_6\text{H}_5\text{COOH}$ . Hence,  $\text{pH}$  is slightly  $> 7$ .

Therefore, increasing order of  $\text{pH}$  of the given salts is,



**Q. 28** The value of  $K_c$  for the reaction  $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$  is  $1 \times 10^{-4}$ . At a given time, the composition of reaction mixture is  $[\text{HI}] = 2 \times 10^{-5}$  mol,  $[\text{H}_2] = 1 \times 10^{-5}$  mol and  $[\text{I}_2] = 1 \times 10^{-5}$  mol. In which direction will the reaction proceed?

**Ans.** Given that,

$$[\text{HI}] = 2 \times 10^{-5} \text{ mol}$$

$$[\text{H}_2] = 1 \times 10^{-5} \text{ mol}$$

$$[\text{I}_2] = 1 \times 10^{-5} \text{ mol}$$

At a given time, the reaction quotient  $Q$  for the reaction will be given by the expression

$$\begin{aligned} Q &= \frac{[\text{H}_2][\text{I}_2]}{[\text{HI}]^2} \\ &= \frac{1 \times 10^{-5} \times 1 \times 10^{-5}}{(2 \times 10^{-5})^2} = \frac{1}{4} \\ &= 0.25 = 2.5 \times 10^{-1} \end{aligned}$$

As the value of reaction quotient is greater than the value of  $K_c$ , i.e.,  $1 \times 10^{-4}$  the reaction will proceed in the reverse reaction.

**Q. 29** On the basis of the equation  $\text{pH} = -\log [\text{H}^+]$ , the  $\text{pH}$  of  $10^{-8} \text{ mol dm}^{-3}$  solution of  $\text{HCl}$  should be 8. However, it is observed to be less than 7.0. Explain the reason.

**Ans.** Concentration  $10^{-8} \text{ mol dm}^{-3}$  indicates that the solution is very dilute. So, we cannot neglect the contribution of  $\text{H}_3\text{O}^+$  ions produced from  $\text{H}_2\text{O}$  in the solution. Total  $[\text{H}_3\text{O}^+] = 10^{-8} + 10^{-7} \text{ M}$ . From this we get the value of  $\text{pH}$  close to 7 but less than 7 because the solution is acidic.

From calculation, it is found that  $\text{pH}$  of  $10^{-8} \text{ mol dm}^{-3}$  solution of  $\text{HCl}$  is equal to 6.96.

**Q. 30**  $\text{pH}$  of a solution of a strong acid is 5.0. What will be the  $\text{pH}$  of the solution obtained after diluting the given solution a 100 times?

**Ans.** Given that,

$$\text{pH} = 5$$

$$[\text{H}^+] = 10^{-5} \text{ mol L}^{-1}$$

$$\text{On diluting the solution 100 times } [\text{H}^+] = \frac{10^{-5}}{100} = 10^{-7} \text{ mol L}^{-1}$$

On calculating the pH using the equation  $\text{pH} = -\log [\text{H}^+]$ , value of pH comes out to be 7. It is not possible. This indicates that solution is very dilute.

Hence, Total  $\text{H}^+$  ion concentration =  $\text{H}^+$  ions from acid +  $\text{H}^+$  ion from water


$$[\text{H}^+] = 10^{-7} + 10^{-7} = 2 \times 10^{-7} \text{ M}$$

$$\text{pH} = -\log [2 \times 10^{-7}]$$

$$\text{pH} = 7 - 0.3010 = 6.699$$

**Q. 31** A sparingly soluble salt gets precipitated only when the product of concentration of its ions in the solution ( $Q_{\text{sp}}$ ) becomes greater than its solubility product. If the solubility of  $\text{BaSO}_4$  in water is  $8 \times 10^{-4} \text{ mol dm}^{-3}$ . Calculate its solubility in  $0.01 \text{ mol dm}^{-3}$  of  $\text{H}_2\text{SO}_4$ .

**Ans.**



$$K_{\text{sp}} \text{ for } \text{BaSO}_4 = [\text{Ba}^{2+}] [\text{SO}_4^{2-}] = s \times s = s^2$$

But

$$s = 8 \times 10^{-4} \text{ mol dm}^{-3}$$

$\therefore$

$$K_{\text{sp}} = (8 \times 10^{-4})^2 = 64 \times 10^{-8}$$

In the presence of  $0.01 \text{ M H}_2\text{SO}_4$ , the expression for  $K_{\text{sp}}$  will be

$$K_{\text{sp}} = [\text{Ba}^{2+}] [\text{SO}_4^{2-}]$$

$$K_{\text{sp}} = (s)(s + 0.01) \quad (0.01 \text{ M } \text{SO}_4^{2-} \text{ ions from } 0.01 \text{ M } \text{H}_2\text{SO}_4)$$

$$64 \times 10^{-8} = s \cdot (s + 0.01)$$

$$s^2 + 0.01s - 64 \times 10^{-8} = 0$$

$$s = \frac{-0.01 \pm \sqrt{(0.01)^2 + (4 \times 64 \times 10^{-8})}}{2}$$

$$= \frac{-0.01 \pm \sqrt{10^{-4} + (256 \times 10^{-8})}}{2}$$

$$= \frac{-0.01 \pm \sqrt{10^{-4}(1 + 256 \times 10^{-4})}}{2}$$

$$= \frac{-0.01 \pm 10^{-2} \sqrt{1 + 0.0256}}{2} = \frac{10^{-2}(-1 \pm 1.012719)}{2}$$

$$= 5 \times 10^{-3}(-1 + 1.012719) = 6.4 \times 10^{-5} \text{ mol dm}^{-3}$$

**Note**  $s \ll 0.01$ , so,  $s + 0.01 \approx 0.01$  and  $64 \times 10^{-8} = s \times 0.01$

$$s = \frac{64 \times 10^{-8}}{0.01} = 6.4 \times 10^{-5}$$

**Q. 32** pH of  $0.08 \text{ mol dm}^{-3}$   $\text{HOCl}$  solution is 2.85. Calculate its ionisation constant.

### 💡 Thinking Process

To solve this problem, we use two steps

**Step I** Find out the concentration of hydrogen ion  $[\text{H}^+]$  through the formula  $-\text{pH} = \log [\text{H}^+]$

**Step II** Afterward, calculate the  $K_a$  of  $\text{HOCl}$  which is weak monobasic acid by using the

formula  $K_a = \frac{[\text{H}^+]^2}{C}$ . where,  $C$  is concentration of the solution

**Ans.**

$$\text{pH of HOCl} = 2.85$$

But,

$$-\text{pH} = \log [\text{H}^+]$$

$\therefore$

$$-2.85 = \log [\text{H}^+]$$

$\Rightarrow$

$$3.15 = \log [\text{H}^+]$$

$\Rightarrow$

$$[\text{H}^+] = 1.413 \times 10^{-3}$$

For weak monobasic acid  $[\text{H}^+] = \sqrt{K_a \times C}$

$\Rightarrow$

$$K_a = \frac{[\text{H}^+]^2}{C} = \frac{(1.413 \times 10^{-3})^2}{0.08} \\ = 24.957 \times 10^{-6} = 2.4957 \times 10^{-5}$$

**Q. 33** Calculate the pH of a solution formed by mixing equal volumes of two solutions A and B of a strong acid having pH = 6 and pH = 4 respectively.

**Ans.** pH of solution A = 6. Hence,  $[\text{H}^+] = 10^{-6} \text{ mol L}^{-1}$

pH of solution B = 4. Hence,  $[\text{H}^+] = 10^{-4} \text{ mol L}^{-1}$

On mixing 1 L of each solution, molar concentration of total  $\text{H}^+$  is halved.

$$\text{Total, } [\text{H}^+] = \frac{10^{-6} + 10^{-4}}{2} \text{ mol L}^{-1}$$

$$[\text{H}^+] = \frac{1.01 \times 10^{-4}}{2} = 5.05 \times 10^{-5} \text{ mol L}^{-1}$$

$$[\text{H}^+] = 5.0 \times 10^{-5} \text{ mol L}^{-1}$$

$$\text{pH} = -\log [\text{H}^+] \quad \Rightarrow \quad \text{pH} = -\log (5.0 \times 10^{-5})$$


$$\text{pH} = -[\log 5 + (-5 \log 10)] \quad \Rightarrow \quad \text{pH} = -\log 5 + 5$$

$$\text{pH} = 5 - \log 5 = 5 - 0.6990 \quad \Rightarrow \quad \text{pH} = 4.3010 \approx 4.3$$

Thus, the pH of resulting solution is 4.3.

**Q. 34** The solubility product of  $\text{Al}(\text{OH})_3$  is  $2.7 \times 10^{-11}$ . Calculate its solubility in  $\text{g L}^{-1}$  and also find out pH of this solution. (Atomic mass of Al = 27 u)

**Ans.** Let S be the solubility of  $\text{Al}(\text{OH})_3$ .



Concentration of species at  $t = 0$

Concentration of various species at equilibrium

$$K_{\text{sp}} = [\text{Al}^{3+}] [\text{OH}^-]^3 = (S) (3S)^3 = 27S^4$$

$$S^4 = \frac{K_{\text{sp}}}{27} = \frac{2.7 \times 10^{-11}}{27} = 1 \times 10^{-12}$$

$$S = 1 \times 10^{-3} \text{ mol L}^{-1}$$

(i) **Solubility of  $\text{Al}(\text{OH})_3$**

Molar mass of  $\text{Al}(\text{OH})_3$  is 78 g. Therefore,

$$\text{Solubility of } \text{Al}(\text{OH})_3 \text{ in } \text{g L}^{-1} = 1 \times 10^{-3} \times 78 \text{ g L}^{-1} = 78 \times 10^{-3} \text{ g L}^{-1} \\ = 7.8 \times 10^{-2} \text{ g L}^{-1}$$

(ii) **pH of the solution**

$$S = 1 \times 10^{-3} \text{ mol L}^{-1}$$

$$[\text{OH}^-] = 3S = 3 \times 1 \times 10^{-3} = 3 \times 10^{-3}$$


$$\text{pOH} = 3 - \log 3$$

$$\text{pH} = 14 - \text{pOH} = 14 + \log 3 = 14.4771$$

**Q. 35** Calculate the volume of water required to dissolve 0.1 g lead (II) chloride to get a saturated solution.

( $K_{sp}$  of  $PbCl_2 = 3.2 \times 10^{-8}$ , atomic mass of  $Pb = 207u$ )

**Ans.** Suppose, solubility of  $PbCl_2$  in water is  $s \text{ mol L}^{-1}$



$$K_{sp} = [s] [2s]^2 = 4s^3$$

$$3.2 \times 10^{-8} = 4s^3$$

$$s^3 = \frac{3.2 \times 10^{-8}}{4} = 0.8 \times 10^{-8}$$

$$s^3 = 8.0 \times 10^{-9}$$

Solubility of  $PbCl_2$ ,  $s = 2 \times 10^{-3} \text{ mol L}^{-1}$

Solubility of  $PbCl_2$  in  $gL^{-1} = 278 \times 2 \times 10^{-3} = 0.556 \text{ g L}^{-1}$

( $\therefore$  Molar mass of  $PbCl_2 = 207 + (2 \times 35.5) = 278$ )


0.556 g of  $PbCl_2$  dissolve in 1 L of water.

$$\therefore 0.1 \text{ g of } PbCl_2 \text{ will dissolve in } = \frac{1 \times 0.1}{0.556} \text{ L of water}$$

$$= 0.1798 \text{ L}$$


To make a saturated solution, dissolution of 0.1g  $PbCl_2$  in 0.1798 L  $\approx$  0.2 L of water will be required.

**Q. 36** A reaction between ammonia and boron trifluoride is given below.



Identify the acid and base in this reaction. Which theory explains it?  
What is the hybridisation of B and N in the reactants?

**Ans.** Although  $BF_3$  does not have a proton but acts as Lewis acid as it is an electron deficient compound. It reacts with  $NH_3$  by accepting the lone pair of electrons from  $NH_3$  and complete its octet. The reaction can be represented by



Lewis electronic theory of acids and bases can explain it. Boron in  $BF_3$  is  $sp^2$  hybridised where N in  $NH_3$  is  $sp^3$  hybridised.

**Q. 37** Following data is given for the reaction  
 $CaCO_3(s) \longrightarrow CaO(s) + CO_2(g)$

$$\Delta_f H^\ominus [CaO(s)] = -635.1 \text{ kJ mol}^{-1}$$

$$\Delta_f H^\ominus [CO_2(g)] = -393.5 \text{ kJ mol}^{-1}$$

$$\Delta_f H^\ominus [CaCO_3(s)] = -1206.9 \text{ kJ mol}^{-1}$$

Predict the effect of temperature on the equilibrium constant of the above reaction.

**Ans.** Given that,

$$\Delta_f H^\ominus [CaO(s)] = -635.1 \text{ kJ mol}^{-1}$$

$$\Delta_f H^\ominus [CO_2(g)] = -393.5 \text{ kJ mol}^{-1}$$

$$\Delta_f H^\ominus [CaCO_3(s)] = -1206.9 \text{ kJ mol}^{-1}$$

In the reaction,



$$\Delta_f H^\ominus = \Delta_f H^\ominus [\text{CaO}(\text{s})] + \Delta_f H^\ominus [\text{CO}_2(\text{g})] - \Delta_f H^\ominus [\text{CaCO}_3(\text{s})]$$

$$\therefore \Delta_f H^\ominus = -635.1 + (-393.5) - (-1206.9) = 178.3 \text{ kJmol}^{-1}$$

Because  $\Delta H$  value is positive, so the reaction is endothermic. Hence, according to Le-Chatelier's principle, reaction will proceed in forward direction on increasing temperature. Thus, the value of equilibrium constant for the reaction increases.

## Matching The Columns

**Q. 38** Match the following equilibria with the corresponding condition.

A. Liquid $\rightleftharpoons$ Vapour	1. Saturated solution
B. Solid $\rightleftharpoons$ Liquid	2. Boiling point
C. Solid $\rightleftharpoons$ Vapour	3. Sublimation point
D. Solute (s) $\rightleftharpoons$ Solute (solution)	4. Melting point
	5. Unsaturated solution

**Ans.** A.  $\rightarrow$  (2)    B.  $\rightarrow$  (4)    C.  $\rightarrow$  (3)    D.  $\rightarrow$  (1)

- A. Liquid  $\rightleftharpoons$  Vapour equilibrium exists at the boiling point.  
 B. Solid  $\rightleftharpoons$  Liquid equilibrium exists at the melting point.  
 C. Solid  $\rightleftharpoons$  Vapour equilibrium exists at the sublimation point.  
 D. Solute (s)  $\rightleftharpoons$  Solute (solution) equilibrium exists at saturated solution.

**Q. 39** For the reaction,  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$

$$\text{Equilibrium constant, } K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2] [\text{H}_2]^3}$$

Some reactions are written below in Column I and their equilibrium constants in terms of  $K_c$  are written in Column II. Match the following reactions with the corresponding equilibrium constant.

Column I (Reaction)	Column II (Equilibrium constant)
A. $2\text{N}_2(\text{g}) + 6\text{H}_2(\text{g}) \rightleftharpoons 4\text{NH}_3(\text{g})$	1. $2K_c$
B. $2\text{NH}_3(\text{g}) \rightleftharpoons 2\text{N}_2(\text{g}) + 3\text{H}_2(\text{g})$	2. $K_c^{1/2}$
C. $\frac{1}{2}\text{N}_2(\text{g}) + \frac{3}{2}\text{H}_2(\text{g}) \rightleftharpoons \text{NH}_3(\text{g})$	3. $\frac{1}{K_c}$
	4. $K_c^2$

**Ans.** A.  $\rightarrow$  (4)    B.  $\rightarrow$  (3)    C.  $\rightarrow$  (2)

For the reaction,  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g})$

$$\text{Equilibrium constant } K_c = \frac{[\text{NH}_3]^2}{[\text{N}_2] [\text{H}_2]^3}$$

A. The given reaction  $[2\text{N}_2(g) + 6\text{H}_2(g) \rightleftharpoons 4\text{NH}_3(g)]$  is twice the above reaction.  
Hence,  $K = K_c^2$

B. The reaction  $[2\text{NH}_3(g) \rightleftharpoons \text{N}_2(g) + 3\text{H}_2(g)]$  is reverse of the above reaction.

Hence,  $K = \frac{1}{K_c}$

C. The reaction  $\left[\frac{1}{2}\text{N}_2(g) + \frac{3}{2}\text{H}_2(g) \rightleftharpoons \text{NH}_3(g)\right]$  is half of the above reaction.

Hence,  $K = \sqrt{K_c} = K_c^{\frac{1}{2}}$ .

**Q. 40** Match standard free energy of the reaction with the corresponding equilibrium constant.

A. $\Delta G^\circ > 0$	1. $K > 1$
B. $\Delta G^\circ < 0$	2. $K = 1$
C. $\Delta G^\circ = 0$	3. $K = 0$
	4. $K < 1$

**Ans.** A.  $\rightarrow (4)$     B.  $\rightarrow (1)$     C.  $\rightarrow (2)$

As we know that,  $\Delta G^\circ = -RT \ln K$

A. If  $\Delta G^\circ > 0$ , i.e.,  $\Delta G^\circ$  is positive, then  $\ln K$  is negative i.e.,  $K < 1$ .

B. If  $\Delta G^\circ < 0$ , i.e.,  $\Delta G^\circ$  is negative then  $\ln K$  is positive i.e.,  $K > 1$ .

C. If  $\Delta G^\circ = 0$ ,  $\ln K = 0$ , i.e.,  $K = 1$ .

**Q. 41** Match the following species with the corresponding conjugate acid.

Species	Conjugate acid
A. $\text{NH}_3$	1. $\text{CO}_3^{2-}$
B. $\text{HCO}_3^-$	2. $\text{NH}_4^+$
C. $\text{H}_2\text{O}$	3. $\text{H}_3\text{O}^+$
D. $\text{HSO}_4^-$	4. $\text{H}_2\text{SO}_4$
	5. $\text{H}_2\text{CO}_3$

**Ans.** A.  $\rightarrow (2)$     B.  $\rightarrow (5)$     C.  $\rightarrow (3)$     D.  $\rightarrow (4)$

As conjugate acid  $\rightarrow$  Base +  $\text{H}^+$




A.  $\text{NH}_3 + \text{H}^+ \longrightarrow \text{NH}_4^+$

B.  $\text{HCO}_3^- + \text{H}^+ \longrightarrow \text{H}_2\text{CO}_3$

C.  $\text{H}_2\text{O} + \text{H}^+ \longrightarrow \text{H}_3\text{O}^+$

D.  $\text{HSO}_4^- + \text{H}^+ \longrightarrow \text{H}_2\text{SO}_4$

**Q. 42** Match the following graphical variation with their description.

<b>A</b>		<b>B</b>
A.		1. Variation in product concentration with time
B.		2. Reaction at equilibrium
C.		3. Variation in reactant concentration with time

**A.** → (3)    **B.** → (1)    **C.** → (2)

A. Graph (A) represents variation of reactant concentration with time.

B. Graph (B) represents variation of product concentration with time.

C. Graph (C) represents reaction at equilibrium.

**Q. 43** Match the Column I with Column II.

<b>Column I</b>	<b>Column II</b>
A. Equilibrium	1. $\Delta G > 0, K < 1$
B. Spontaneous reaction	2. $\Delta G = 0$
C. Non-spontaneous reaction	3. $\Delta G^\ominus = 0$
	4. $\Delta G < 0, K > 1$

**Ans.** **A.** → (2, 3)    **B.** → (4)    **C.** → (1)

A.  $\Delta G$  ( $\Delta G^\ominus$ ) is 0, reaction has achieved equilibrium: at this point, there is no longer any free energy left to drive the reaction.

B. If  $\Delta G < 0$ , then  $K > 1$  which implies a spontaneous reaction or the reaction which proceeds in the forward direction to such an extent that the products are present predominantly.

C. If  $\Delta G > 0$ , then  $K < 1$ , which implies a non-spontaneous reaction or a reaction which proceeds in the forward direction to such a small degree that only a very minute quantity of product is formed.

## Assertion and Reason

In the following questions a statement of Assertion (A) followed by a statement of Reason (R) is given. Choose the correct option out of the choices given below in each question.

**Q. 44 Assertion (A)** Increasing order of acidity of hydrogen halides is  $\text{HF} < \text{HCl} < \text{HBr} < \text{HI}$ .

**Reason (R)** While comparing acids formed by the elements belonging to the same group of periodic table,  $\text{H}-\text{A}$  bond strength is a more important factor in determining acidity of an acid than the polar nature of the bond.

- (a) Both A and R are true R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is true but R is false
- (d) Both A and R are false

**Ans. (a)** Both assertion and reason are true and reason is the correct explanation of assertion.

In the hydrogen halides, the HI is strongest acid while HF is the weak acid. It is because while comparing acids formed by the elements belonging to the same group of periodic table,  $\text{H}-\text{A}$  bond strength is a more important factor in determining acidity of an acid than the polar nature of the bond.

**Q. 45 Assertion (A)** A solution containing a mixture of acetic acid and sodium acetate maintains a constant value of pH on addition of small amounts of acid or alkali.

**Reason (R)** A solution containing a mixture of acetic acid and sodium acetate acts as buffer solution around pH 4.75.

- (a) Both A and R are true and R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is true but R is false
- (d) Both A and R are false

**Ans. (a)** Both assertion and reason are true and reason is correct explanation of assertion.

A solution containing a mixture of acetic acid and the sodium acetate acts as a buffer solution as it maintains a constant value of pH ( $= 4.75$ ) and its pH is not affected on addition of small amounts of acid or alkali.

**Q. 46 Assertion (A)** The ionisation of hydrogen sulphide in water is low in the presence of hydrochloric acid.

**Reason (R)** Hydrogen sulphide is a weak acid.

- (a) Both A and R are true and R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is true but R is false
- (d) Both A and R are false

**Ans. (b)** Both assertion and reason are true but reason is not correct explanation of assertion.

$\text{HCl}$  gives the common  $\text{H}^+$  ions and hence ionisation equilibrium  $\text{H}_2\text{S} \rightleftharpoons 2\text{H}^+ + \text{S}^{2-}$  is suppressed.

**Q. 47 Assertion (A)** For any chemical reaction at a particular temperature, the equilibrium constant is fixed and is a characteristic property.

**Reason (R)** Equilibrium constant is independent of temperature.

- (a) Both A and R are true and R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is true but R is false
- (d) Both A and R are false

**Ans. (c)** Assertion is true but reason is false.


Equilibrium constant of a reaction depends upon temperature.

**Q. 48 Assertion (A)** Aqueous solution of ammonium carbonate is basic.

**Reason (R)** Acidic/basic nature of a salt solution of a salt of weak acid and weak base depends on  $K_a$  and  $K_b$  value of the acid and the base forming it.

- (a) Both A and R are true and R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is true but R is false
- (d) Both A and R are false

**Ans. (a)** Both assertion and reason are true and reason is the correct explanation of assertion.



If  $K_b$  of  $\text{NH}_4\text{OH} > K_a$  of  $\text{H}_2\text{CO}_3$

The solution is basic.

or, if  $K_a$  of  $\text{H}_2\text{CO}_3 > K_b$  of  $\text{NH}_4\text{OH}$ ; the solution is acidic.

**Q. 49 Assertion (A)** An aqueous solution of ammonium acetate can act as a buffer.

**Reason (R)** Acetic acid is a weak acid and  $\text{NH}_4\text{OH}$  is a weak base.

- (a) Both A and R are true and R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is false but R is true
- (d) Both A and R are false

**Ans. (b)** Both assertion and reason are true but reason is not correct explanation of assertion.


Ammonium acetate is a salt of weak acid ( $\text{CH}_3\text{COOH}$ ) and weak base ( $\text{NH}_4\text{OH}$ ).

**Q. 50 Assertion (A)** In the dissociation of  $\text{PCl}_5$  at constant pressure and temperature the addition of helium at equilibrium increases the dissociation of  $\text{PCl}_5$ .

**Reason (R)** Helium removes  $\text{Cl}_2$  from the field of action.

- (a) Both A and R are true and R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is true but R is false
- (d) Both A and R are false

**Ans. (c)** Assertion is true but reason is false.



At constant pressure, when helium is added to the equilibrium, volume increases. Thus, in order to maintain the  $K$  constant, degree of dissociation of  $\text{PCl}_5$  increases. Helium is unreactive towards chlorine gas.

## Long Answer Type Questions

**Q. 51** How can you predict the following stages of a reaction by comparing the value of  $K_c$  and  $Q_c$ ?

- (i) Net reaction proceeds in the forward direction.
- (ii) Net reaction proceeds in the backward direction.
- (iii) No net reaction occurs.

**Ans.** Prediction of the following stages of a reaction by comparing the value of  $K_c$  and  $Q_c$  are

- (i) If  $Q_c < K_c$ , the reaction will proceed in the direction of the products (forward reaction).
- (ii) If  $Q_c > K_c$ , the reaction will proceed in the direction of reactants (reverse reaction).
- (iii) If  $Q_c = K_c$ , the reaction mixture is already at equilibrium.

**Q. 52** On the basis of Le-Chatelier principle explain how temperature and pressure can be adjusted to increase the yield of ammonia in the following reaction.



What will be the effect of addition of argon to the above reaction mixture at constant volume?

**Ans.**  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightleftharpoons 2\text{NH}_3(\text{g}); \Delta H = -92.38 \text{ kJ mol}^{-1}$

It is an exothermic process as  $\Delta H$  is negative.

**Effect of temperature** According to Le-Chatelier's principle, low temperature is favourable for high yield of ammonia, but practically very low temperatures slow down the reaction. So, optimum temperature 700 K is favourable in attainment of equilibrium.

**Effect of pressure** Similarly, high pressure about 200 atm is favourable for high yield of ammonia. On increasing pressure, reaction goes in the forward direction because the number of moles decreases in the forward direction.

**Addition of argon** At constant volume addition of argon does not affect the equilibrium because it does not change the partial pressures of the reactants or products involved in the reaction and the equilibrium remains undisturbed.

**Q. 53** A sparingly soluble salt having general formula  $A_x^{p+} B_y^{q-}$  and molar solubility  $S$  is in equilibrium with its saturated solution. Derive a relationship between the solubility and solubility product for such salt.

**Ans.** A sparingly soluble salt having general formula  $A_x^{p+} B_y^{q-}$ . Its molar solubility is  $S \text{ mol L}^{-1}$ .

Then,  $A_x^{p+} B_y^{q-} \rightleftharpoons x A_x^{p+} (aq) + y B_y^{q-} (aq)$

$S$  moles of  $A_x B_y$  dissolve to give  $x$  moles of  $A^{p+}$  and  $y$  moles of  $B^{q-}$ .

Therefore, solubility product ( $K_{sp}$ ) =  $[A^{p+}]^x [B^{q-}]^y$

$$= [xS]^x [yS]^y$$

$$= x^x y^y S^{x+y}$$

**Q. 54** Write a relation between  $\Delta G$  and  $Q$  and define the meaning of each term and answer the following.

(a) Why a reaction proceeds forward when  $Q < K$  and no net reaction occurs when  $Q = K$ ?

(b) Explain the effect of increase in pressure in terms of reaction quotient  $Q$ .

For the reaction,  $\text{CO(g)} + 3\text{H}_2\text{(g)} \rightleftharpoons \text{CH}_4\text{(g)} + \text{H}_2\text{O(g)}$

**Ans.** The relation between  $\Delta G$  and  $Q$  is

$$\Delta G = \Delta G^\circ + RT \ln Q$$

$\Delta G$  = change in free energy as the reaction proceeds.

$\Delta G^\circ$  = standard free energy

$Q$  = reaction quotient

$R$  = gas constant

$T$  = absolute temperature in  $K$

(a) Since,

$$\Delta G^\circ = -RT \ln K$$

$\therefore$

$$\Delta G = -RT \ln K + RT \ln Q$$

$$\Delta G = RT \ln \frac{Q}{K}$$

If  $Q < K$ ,  $\Delta G$  will be negative and the reaction proceeds in the forward direction.

If  $Q = K$ ,  $\Delta G = 0$  reaction is in equilibrium and there is no net reaction.

(b)  $\text{CO(g)} + 3\text{H}_2\text{(g)} \rightleftharpoons \text{CH}_4\text{(g)} + \text{H}_2\text{O(g)}$

$$K_c = \frac{[\text{CH}_4] [\text{H}_2\text{O}]}{[\text{CO}] [\text{H}_2]^3}$$

On increasing pressure, volume decreases. If we doubled the pressure, volume will be halved but the molar concentrations will be doubled. Then,

$$Q_c = \frac{2[\text{CH}_4] \cdot 2[\text{H}_2\text{O}]}{2[\text{CO}] \{2[\text{H}_2]\}^3} = \frac{1}{4} \frac{[\text{CH}_4] [\text{H}_2\text{O}]}{[\text{CO}] [\text{H}_2]^3} = \frac{1}{4} K_c$$

Therefore,  $Q_c$  is less than  $K_c$ , so  $Q_c$  will tend to increase to re-establish equilibrium and the reaction will go in forward direction.