LOCOMOTION AND MOVEMENT

Locomotion is the voluntary movements resulting in a in location. All locomotion are movements but all movements are not locomotion. Both are interlinked. E.g.

- In *Paramoecium*, cilia help in the movement of food through cytopharynx and in locomotion.
- *Hydra* use tentacles to capture prey and for locomotion.
- Limbs help to change body postures and for locomotion.

Types of movement in human being

• Amoeboid movement: By pseudopodia formed by streaming of protoplasm as in *Amoeba*. Cytoskeletal


HUMAN MUSCULAR SYSTEM

- It includes muscles which are mesodermal in origin.
- Muscles constitute 40-50% of the body weight.
- Muscles have excitability, contractility, extensibility & elasticity.
- Based on location, muscles are 3 types:

Skeletal (striated) muscles	Visceral (Non- striated) muscles	Cardiac muscles
Attached to skeleton	In visceral organs	In heart wall
Striations present	Absent	Present
Voluntary	Involuntary	Involuntary
Rich blood supply	Poor blood supply	Rich blood supply
Fatigue muscle	Non-fatigue	Non-fatigue
Multinucleate	Uninucleate	Uninucleate
More mitochondria	Less mitochondria	More mitochondria

STRUCTURE OF STRIATED MUSCLE

- Skeletal muscle is made of **muscle bundles (fascicles)** held together by collagenous connective tissue layer **(fascia)**.





- Each fascicle contains many muscle fibres (muscle cells).
- Muscle fibres are lined by **plasma membrane** (sarcolemma) enclosing the sarcoplasm.
- Each muscle fibre contains myofilaments (myofibrils).
- Each myofibril has alternate dark (Anisotropic or Aband) and light striations (Isotropic or I-band). This is due to the presence of 2 fibrous contractile proteins- thin Actin filament and thick Myosin filament.
- I-bands contain actin. A-bands contain actin and myosin. They are arranged parallel to each other.
- A-band bears a lighter middle region (**H band**) formed of only myosin. A thin dark line (**M-line**) runs through the centre of **H-zone**.
- I-band is bisected by a dense dark band called **Z-line**. Region between two Z-lines is called **sarcomere**. They are the *functional units of muscle contraction*.

elements like microfilaments also help for this. E.g. change Macrophages & leucocytes.

- Ciliary movement: By cilia. E.g. ciliary movements in trachea (to remove dust particles and foreign substances), and oviducts (for the passage of ova).
- Muscular movement: By muscles. E.g. movement of limbs.


Flagellar movement helps in the swimming of spermatozoa, maintenance of water current in the canal system of sponges and in locomotion of Protozoans like *Euglena*.






Structure of contractile proteins

- An actin filament is made of 2 filamentous (F) actins which form double helix.
- F-actin is a polymer of monomeric Globular (G) actins.



- Actin contains 2 other proteins (tropomyosin & troponin).
- Two filaments of **tropomyosin** run along the grooves of the F-actin double helix.
- **Troponin** has 3 subunits. It is seen at regular intervals on tropomyosin. In the resting state, **a subunit of troponin** masks the binding sites for myosin on the actin filaments.
- Each myosin filament is a polymer of many **monomeric proteins** called **Meromyosins**.



- A meromyosin has 2 parts:
 - Heavy meromyosin or HMM or cross arm (globular head + short arm): It projects outwards.
 - Light meromyosin or LMM (tail).
- The globular head is an active *ATPase* enzyme and has binding sites for ATP and active sites for actin.

MECHANISM OF MUSCLE CONTRACTION


According to sliding filament theory, contraction of a muscle fibre occurs by the sliding of thin filaments over thick filaments. The steps are given below:

• An impulse from the CNS reaches the neuromuscular junction (Motor-end plate) via motor neuron.

Neuromuscular junction is the synapse between a motor neuron and the sarcolemma of the muscle fibre.

A motor neuron + muscle fibres = a motor unit.

- Synaptic vesicles release a neurotransmitter Acetylcholine. It generates an **action potential** in the sarcolemma that spreads through the muscle fibre. It causes the release of Ca²⁺ ions from sarcoplasmic cisternae into sarcoplasm.
- Ca binds with a subunit of troponin on actin filaments and unmask the active sites for myosin.



Using energy from ATP hydrolysis, myosin head binds to active sites on the actin to form cross bridge. This pulls actin filaments on both sides towards the centre of A-band. Actin filaments partially overlap so that H-zone disappears.


It consists of a framework of bones (206) & few cartilages. Human skeletal system has 2 parts: axial & appendicular.

1. Axial skeletal system (80 bones)

Includes bones of head, vertebral column, sternum & ribs.


a. Bones of Head (29 bones)

It includes skull, Hyoid and Ear ossicles.



• Skull (22): Include cranial bones and facial bones. Cranial bones (8): Include Frontal (1), Parietals (2), Temporals (2), Occipital (1), Sphenoid (1) & Ethmoid (1). Facial bones (14): Include Nasals (2), Maxillae (2), Zygomatics (2), Lacrimals (2), Palatines (2), Inferior nasals (2), Mandible (1) and Vomer (1).

- The Z- line attached to actins is also pulled inwards. It causes a shortening (contraction) of sarcomere.
- I-bands get shortened, whereas A-bands retain the length.
- Myosin releases ADP and Pi and goes back to its relaxed state. A new ATP binds and the cross-bridge is broken.
- The ATP is again hydrolyzed by the myosin head and the above processes are repeated causing further sliding.
- When Ca²⁺ ions are pumped back to sarcoplasmic cisternae, actin filaments are again masked. As a result, Z-lines return to their original position. It results in relaxation.



- The reaction time of the fibres varies in different muscles.
- Repeated activation of muscles leads to the accumulation of the lactic acid causing muscle fatigue. This is due to anaerobic breakdown of glycogen in muscles.

Red muscle fibres and white muscle fibres


Red (Aerobic) muscles	White muscle
Red coloured due to myoglobin	White coloured due to lesser myoglobin
More mitochondria	Less mitochondria
Aerobic metabolism	Anaerobic metabolism
Slow & sustained contraction	Fast contraction for short period

HUMAN SKELETAL SYSTEM Skull articulates with First vertebra (atlas) with the help of 2 occipital condyles (dicondylic skull).


- Hyoid bone (1): U-shaped bone seen below buccal cavity.
- Ear ossicles $(3 \times 2 = 6)$: Malleus (2), Incus (2) & stapes (2).

b. Vertebral column

- Formed of **26 vertebrae**. Includes *Cervical vertebrae* (7). Thoracic vertebrae (12), Lumbar vertebrae (5), Sacral vertebrae (1-fused) and Coccygeal vertebrae (1-fused).
- Vertebra has a central hollow portion (neural canal) through which the spinal cord passes.
- Number of cervical vertebrae are 7 in almost all mammals.
- The vertebral column protects the spinal cord, supports the head and serves as the point of attachment for the ribs and musculature of the back.




- Flat bone on the ventral midline of thorax.



- **True ribs** (first 7 pairs): They are attached to *thoracic vertebrae* and ventrally connected to sternum with the help of *Hyaline cartilage*.
- Vertebrochondral (false) ribs (8th, 9th & 10th pairs): They do not articulate directly with the sternum but join the 7^{th} *rib* with the help of *Hyaline cartilage*.
- **Floating ribs** (11th & 12th pairs): They are not connected ventrally (no connection with sternum or other ribs).
- Each rib has 2 articulation surfaces on its dorsal end and is hence called **bicephalic.**

2. Appendicular skeletal system (126 bones)



a. Bones of fore-limbs (30 x 2 = 60)

Include Humerus (1), Radius (1), Ulna (1), Carpals (wrist bones-8), Metacarpals (palm bones-5) & Phalanges (digits-14).

DISORDERS OF MUSCULAR & SKELETAL SYSTEMS

- **Myasthenia gravis:** An auto immune disorder that affects neuromuscular junction. It leads to fatigue, weakening and paralysis of skeletal muscles.
- **Muscular dystrophy:** Progressive degeneration of skeletal muscles. Mostly due to genetic disorder.
- Tetany: Rapid muscle spasm due to low Ca²⁺ in body fluid.

b. Bones of hind-limbs (30 x 2 = 60)

Include Femur (thigh bone- 1), Patella (knee cap- 1), Tibia (1) & fibula (1), Tarsals (ankle bones-7), Metatarsals (5) & Phalanges (digits-14).

c. Pectoral girdles (2x2=4)

- Include clavicle (2) & scapula (2).
- Scapula is a large *triangular* flat bone situated in the dorsal part of the thorax between *the second and* 7th *ribs*.
- Scapula (shoulder blade) has a slightly elevated ridge (*spine*) which projects as a flat, expanded process (*acromion*). The *clavicle* (collarbone) articulates with this.
- Below the acromion is *glenoid cavity* which articulates with the head of *humerus* to form the *shoulder joint*.

d. Pelvic girdles (2)

- Formed of 2 coxal bones. Each coxal bone is formed by the fusion of 3 bones- *Ilium, Ischium & pubis.*
- At the point of fusion of *Ilium*, *Ischium* and *Pubis* is a cavity *(Acetabulum)* to which the *thigh bone* articulates.
- The 2 halves of the *pelvic girdle* meet ventrally to form *pubic symphisis* containing *fibrous cartilage*.

JOINTS

Joints are points of contact between bones, or between bones and cartilages. 3 types:

- 1. Fibrous (immovable) joints: E.g. sutures b/w skull bones.
- 2. Cartilaginous joints (Slightly movable joints): Bones are joined together with the help of cartilages. E.g. Joints between the adjacent vertebrae.
- **3.** Synovial (movable) joints: They have a fluid filled synovial cavity between articulating surfaces of 2 bones.

Types of synovial joint

Joint	Examples	
Ball & socket	Shoulder joint & hip joints.	
Hinge joint	Knee joint, elbow joint, phalanges joints	
Pivot joint	Joints b/w atlas & axis.	
Gliding joint	Joints b/w carpals	
Saddle joint	Joints b/w carpal & metacarpal of thumb	

- Arthritis: Inflammation of joints.
- Osteoporosis: Age-related disorder characterized by decreased bone mass and increased chances of fractures. Decreased level of estrogen is a common cause.
- Gout: Inflammation of joints due to accumulation of uric acid crystals.

